

OTTO FILE CHOS



0-A201 79

Institute Report No. 296

Mutagenic Potential of Triethyleneglycol Dinitrate in the Ames Salmonella/Mammalian Microsome Mutagenicity Test



Steven K. Sano, BA, SGT, USA and Don W. Korte, Jr., PhD, MAJ, MSC

GENETIC TOXICOLOGY BRANCH
DIVISION OF TOXICOLOGY

DISTRIBUTION STATEMENT A

Approved for public releases
Distribution Unlimited

September 1988

Toxicology Series: 113

LETTERMAN ARMY INSTITUTE OF RESEARCH PRESIDIO OF SAN FRANCISCO, CALIFORNIA 94129

This document has been approved for public release and sale; its distribution is unlimited.

Destroy this report when it is no longer needed. Do not return to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360-5)

Edwin S. Beatrice

COL, MC Commanding 28 July 88 (date)

UNCLASSIFIED SECURITY CLASSIFICATION OF THIS PAGE

REPORT	DOCUMENTATIO	N PAGE			Form Approved OMB No. 0704-0188
1a. REPORT SECURITY CLASSIFICATION		1b. RESTRICTIVE	MARKINGS		
Unclassified 2a. SECURITY CLASSIFICATION AUTHORITY		2 DISTRIBUTION	(AVAII ABII ITS		
		3. DISTRIBUTION Approved f	or public	OF REPORT release:	distribution
2b. DECLASSIFICATION/DOWNGRADING SCHED	ULE	is unlim	ited.		- Alleys
4. PERFORMING ORGANIZATION REPORT NUMB	ER(S)	5. MONITORING	ORGANIZATION	REPORT NU	MBER(S)
Institute Report No. 296					ere des
6a. NAME OF PERFORMING ORGANIZATION Genetic Toxicology Branch	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MO			
Division of Toxicology	SGRD-ULE-T	Laborator	eaicai kes	earch ar	nd Development
6c. ADDRESS (City, State, and 7IP Code)		7b. ADDRESS (Ch		Code)	
Letterman Army Institute of Res	search	Ft. Detrick			
Presidio of San Francisco, CA	94129-6800	Frederick,	MD 21701-5	010	
Ba. NAME OF FUNDING/SPONSORING	8b. OFFICE SYMBOL	9. PROCUREMENT	I INSTRUMENT I	DENTIFICAT	ION NUMBER
ORGANIZATION US Army Medical Research & Development Command	(If applicable)				:
8c. ADDRESS (City, State, and ZIP Code)	1	10. SOURCE OF F		.04	
Ft. Detrick		PROGRAM	PROJECT	TASK	WORK UNIT
Frederick, MD 21701-5012		ELEMENT NO.	NO.	NO.	ACCESSION NO.
	· 	62720A	835	AB	DA 303913
11. TITLE (Include Security Classification) Muta SalmoneIla/Mammalian Microsome 12. PERSONAL AUTHOR(S) Steven K. Sano and Don W. I	(orte, Jr.				
والمستبين والمتقود والأوار والمناف والمناف والمناف والمناف والمناف والمناف والمناف والمناف والمناف والماف والم	COVERED 17/85 to 1/25/85	14. DATE OF REPO 1988	RT (<i>Year, Mont!</i> September	n, Day) 15	. PAGE COUNT 20
16. SUPPLEMENTARY NOTATION	Toxicology Ser	ies 113			
17. COSATI CODES	18. SUBJECT TERMS (-	_	=
FIELD GROUP SUB-GROUP	Mutagenicity / Genetic toxico	TRIETH	YLENEGLYCO	L DINIT	RATE
	Ames Test,	logy/ TEGDN/ Propel		75.25	
19 ABSTRACT (Continue on reverse if necessar			·		
The mutagenic potential of Salmonella/Mammalian Microsome TAI537, and TAI538 were exposed the presence and absence of met conditions of this test.	Mutagenicity Tes I to doses rangin tabolic activatio	t. Tester s g from 5 №2/	trains TA9 plate to 0 compound	8, TA100 .0016 ⊾	\ ''TATET "'
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT		21. ABSTRACT SE		CATION	
MANG OF RECOVERS AND WOULD	RPT. DTIC USERS	unclassi		w 1 3 5 - A 5	erce evanes:
22a. NAME OF RESPONSIBLE INDIVIDUAL Edwin S. Beatrice, COL, M	IC .	415-561-			P-UL-Z
DD Form 1473, JUN 86	Previous editions are				ATION OF THIS PAGE

ABSTRACT

The mutagenic potential of TRIETHYLENEGLYCOL DINITRATE was assessed by using the Ames <code>Salmonella/Mammalian</code> Microsome Mutagenicity Test. Tester strains TA98, TA100, TA1535, TA1537, and TA1538 were exposed to doses ranging from 5 μ l/plate to 0.0016 μ l/plate in both the presence and absence of metabolic activation. The test compound was not mutagenic under conditions of this test.

Key Words: Mutagenicity, Genetic Toxicology, Ames Test, TRIETHYLENEGLYCOL DINITRATE, TEGDN, Propellant



Acces	iOt, i r	·
NTIS DTIC	Official Tells	V
	omi i	
By Distrib		
.4	walling of	· · · · · · · · · · · · · · · · · · ·
Dist	Association (
A-1		

PREFACE

TYPE REPORT: Ames Test GLP Study Report

TESTING FACILITY:

US Army Medical Research and Development Command Letterman Army Institute of Research Presidio of San Francisco, CA 94129-6800

SPONSOR:

US Army Medical Research and Development Command
US Army Biomedical Research and Development Laboratory
Fort Detrick, MD 21701-5010
Project Officer: Gunda Reddy, PhD

PROJECT/WORK UNIT/APC: 3E162720A835/180/TLBO

GLP STUDY NUMBER: 84048

STUDY DIRECTOR: MAJ Don W. Korte Jr, PhD, MSC

PRINCIPAL INVESTIGATOR: Steven K. Sano, BA, SGT, USA

REPORT AND DATA MANAGEMENT:

A copy of the final report, study protocol, retired SOPs, stability, and purity data of the test compound, and an aliquot of the test compound will be retained in the LAIR Archives.

TEST SUBSTANCE: TRIETHYLENEGLYCOL DINITRATE (TEGDN)

INCLUSIVE STUDY DATES: 7-25 January 1985

OBJECTIVE:

The objective of this study was to determine the mutagenic potential of TRIETHYLENEGLYCOL DINITRATE (LAIR Code TA044) by using the Ames Salmonella/Mammalian Microsome Mutagenicity Test.

ACKNOWLEDGMENTS

CPT John W. Harbell, PhD, MSC, and Mr. John Dacey provided research assistance.

SIGNATURES OF PRINCIPAL SCIENTISTS INVOLVED IN THE STUDY

We, the undersigned, declare that GLP study number 84048 was performed under our supervision, according to the procedures described herein, and that this report is an accurate record of the results obtained.

OON W. KORTE JR. PHD// DATE

MAJ, MSC

Study Director

CONNAD WHEELER PADY DATE

DAC

Analytical chemist

Steven Karuo Sano 5 MAR 86

STEVEN K. SANG/, BA / DATE

SGT, USA

Principal Investigator



DEPARTMENT OF THE ARMY

LETTERMAN ARMY INSTITUTE OF RESEARCH PRESIDIO OF SAN FRANCISCO CALIFORNIA 94129-6800

REPENTON OF

SGRD-ULZ-QA

22 September 1988

MEMORANDUM FOR RECORD

SUBJECT: GLP Compliance Statement

- 1. This is to certify that the protocol for GLP Study 84048 was reviewed on 4 December 1984.
- 2. The institute report entitled "Mutagenic Potential of Triethyleneglycol Dinitrate in the Ames Salmonella/Mammalian Microsome Mutagenicity Test," Toxicology Series 113, was audited on 12 May 1987.

Carolyn M. LEWIS

Chief, Quality Assurance

TABLE OF CONTENTS

PAGE
Abstract i
Prefaceiii
Acknowledgmentsiv
Signatures of Principal Scientistsv
Report of the Quality Assurance Unitvi
Table of Contentsvii
INTRODUCTION 1
Objective of the Study1
MATERIALS AND METHODS
Test Compound 2
Test Solvent 2
Chemical Preparation 2
Test Strains 2
Test Format 3
Data Interpretation 4
Deviations from the Protocol/SOP 4
Storage of Raw Data and Final Report 4
RESULTS 5
DISCUSSION 5
CONCLUSION
PEREDENOES 11

TABLE OF CONTENTS

PAGE	3
APPENDICES	
APPENDIX A	
APPENDIX B	
OFFICIAL DISTRIBUTION LIST	

Mutagenic Potential of TRIETHYLENEGLYCOL DINITRATE in the Ames Salmonella/Mammalian Microsome Mutagenicity Test--Sano and Korte

INTRODUCTION

The Department of Defense is considering the use of diethyleneglycol dinitrate (DEGDN), triethyleneglycol dinitrate (TEGDN), or trimethylolethane trinitrate (TMETN) as a replacement for nitroglycerin in munition formulations. A "health effects" review conducted for the US Army Biomedical Research and Development Laboratory (USABRDL) identified numerous gaps in the toxicology database of these compounds (1). Consequently, USABRDL has tasked the Toxicology Branch, LAIR, to conduct an initial health effects evaluation of DEGDN, TMETN, TEGDN, and two DEGDN-based propellants, JA-2 and DIGL-RP. This initial evaluation includes the Ames mutagenicity test, acute oral toxicity tests in rats and mice, acute dermal toxicity tests in rabbits, dermal and ocular irritation studies in rabbits, and dermal sensitization studies in quinea pigs. This report contains the results of a study to assess the mutagenic potential of TEGDN in the Ames Salmonella/Mammalian Microsome Mutagenicity Test.

The Ames Salmonella/Mammalian Microsome Mutagenicity Test is a short-term screening test that utilizes histidine auxotrophic mutant strains of Salmonella typhimurium to detect those compounds which are potentially mutagenic in mammals. A mammalian microsomal enzyme system is incorporated in the test to increase sensitivity by simulating in vivo metabolic activation of the test compound. The Ames test is an inexpensive yet highly predictive and reliable test for detecting mutagenic activity and thus carcinogenic potential (2).

Objective of the Study

The objective of this study was to determine the mutagenic potential of TRIETHYLENEGLYCOL DINITRATE (LAIR Code TA044) by using the Ames Salmonella/Mammalian Microsome Mutagenicity Test.

Sano and Korte--2

MATERIAL AND METHODS

Test Compound

Chemical Name: TRIETHYLENEGLYCOL DINITRATE

LAIR Code Number: TA044

Physical State: yellow oil

Source: Naval Ordnance Station

Indian Head, MD

Storage: TRIETHYLENEGLYCOL DINITRATE was received and assigned the LAIR Code number TA044. The test compound was stored at room temperature (21°C) until use.

Chemical Properties/Analysis: Data characterizing the chemical composition and purity of the test material are presented in Appendix A.

Test Solvent

The test compound and the positive control chemicals were dissolved in grade 1 dimethyl sulfoxide (lot 113F-0450) obtained from Sigma Chemical Co. (St. Louis, MO).

Chemical Preparation

On the day before dosing, 500 μ l of the test compound was measured into a sterile vial and again stored at room temperature. On the day of dosing, the 500- μ l sample was dissolved in 9.4 ml of grade I dimethyl sulfoxide to achieve a 5% (v/v) solution. Aliquots of this solution were used to dose the test plates.

Test Strains

Salmonella strains TA98, TA100, TA1535, TA1537, and TA1538, obtained directly from Dr. Bruce Ames, University of California, Berkeley, were used. These strains were maintained in our laboratory at -80°C. Quality control tests were run concurrently with the test substance to establish the validity of their special features and to determine the spontaneous reversion rate. Descriptions of the strains, their genetic markers, and the methods for strain validation are given in the LAIR SOP, OP-STX-1 (3).

Test Format

TRIETHYLENEGLYCOL DINITRATE was evaluated for mutagenic potential according to the methods of Ames et al (4). A detailed description of the methodology is given in LAIR SOP, OP-STX-1 (3).

Toxicity Tests: Toxicity tests were conducted to determine a sublethal concentration of the test substance. This toxicity level was found by using minimal glucose agar (MGA) plates, concentrations of TRIETHYLENEGLYCOL DINITRATE ranging from 1.6 x 10^{-3} µl/plate to 5 µl/plate and approximately 10^8 cells of TA100 per plate. Top agar containing trace amounts of histidine and biotin was placed on the plates. Strain verification was confirmed on the bacteria, along with a determination of the spontaneous reversion rate.

Mutagenicity Test: The test substance was evaluated over a 1000-fold range of concentrations, decreasing from the minimum toxic level (the maximum or limit dose) by a dilution factor of 5, both with and without 0.5 ml of the S-9 microsome fraction. The S-9 was purchased from Litton Bionetics (Kensington, MD). The optimal titer of this S-9, as determined by Litton Bionetics, was 0.75 mg protein/plate. After all the ingredients were added, the top agar was mixed, then overlaid on MGA plates. These plates contained 2% glucose and Vogel Bonner "E" concentrate (5). The water used in this medium and in all reagents came from a Polymetric model 200-3 Water Purifier (Sunnyvale, CA). Plates were incubated upside down in the dark, at 37°C for 48 hours. Plates were prepared in triplicate and the average revertant counts were recorded. The average number of revertants at each dose level was compared to the average number of spontaneous revertants (negative control). The spontaneous reversion rate (with and without S-9) was monitored by averaging the counts from two determinations run simultaneously with the test compound. The spontaneous reversion rate was determined by inoculating one set of plates before and one set after the test compound plates so that any change in spontaneous reversion rate during the dosing procedure would be detected. This spontaneous reversion rate was also compared with historical values for this laboratory and those cited in Ames et al (4). Sterility and strain verification controls were run concurrently. All reagents, test compounds, and media were checked for sterility by plating samples of each on MGA media and incubating them at 37°C with the test plates. The integrity

of the different Salmonella strains used in the assay was verified by the following standard tests:

-Lack of growth (inhibition) in the presence of crystal violet which indicates that the prerequisite alteration of the lipopolysaccharide layer of the cell wall is present.

-Growth in the presence of ampicillin-impregnated disks which indicates the presence of an ampicillin-resistant R Factor in the TA98 and TA100 strains.

-Lack of growth (inhibition) following exposure to ultraviolet light indicates the absence of the DNA excision-repair mechanism.

Four known mutagens were tested as positive controls to confirm the responsiveness of the strains to the mutation process. Each strain must be tested with at least one positive control, but may be tested with several. These compounds (benzo[a]pyrene, 2-aminofluorene, 2-aminoanthracene and N-methyl-N'-nitro-N-nitrosoguanidine) were obtained from Sigma Chemical Co. (St. Louis, MO). The test compound and the known mutagens were handled during this study in accordance with the standards published in NIH Guidelines for the Laboratory Use of Chemical Carcinogens (DHHS Publication No. (NIH) 81-2385, May 1981).

Data Interpretation

According to Brusick (6), a compound is considered mutagenic if a positive dose response (correlated dose response) over three dose concentrations is achieved with at least the highest dose yielding a revertant colony count for the tester strains TA98 and TA100, or three times the spontaneous colony count for strains TA1535, TA1537, and TA1538. A strong correlated dose response in strain TA100 without a doubling of the individual colony count may also be considered positive.

Deviations from the Protocol/SOP

None.

Storage of Raw Data and Final Report

A copy of the final report, study protocols, raw data, SOPs, and an aliquot of the test compound will be retained in the LAIR archives.

RESULTS

On 16 January 1985, the toxicity of TRIETHYLENEGLYCOL DINITRATE was determined (Table 1). For this experiment all sterility, strain verification, and negative controls were normal (Table 1). Exposure of the tester strain (TA100) to the highest dose showed neither a decrease in the number of macrocolonies nor an observable reduction in the density of the background lawn, indicating no chemical toxicity. Therefore, the highest dose selected for the mutagenicity test was $5~\mu l/plate$.

Normal results were obtained for all sterility and strain verification tests during the Ames Test performed on 23-25 January 1985 (Table 2). TRIETHYLENEGLYCOL DINITRATE did not induce any appreciable increase in the revertant colony counts relative to those of the negative control cultures (Table 3).

A copy of the raw data is included in Appendix B.

DISCUSSION

Certain test criteria must be satisfied before the Ames Test can be considered a valid assessment of a compound's mutagenic potential. First, the special features of the Ames strains must be verified. These features include demonstration of ampicillin resistance, alteration in the LP (lipopolysaccharide) layer, and deficiency in DNA excision-repair. Second, the Salmonella strains must be susceptible to mutation by known mutagens. Third, the optimal concentration of the test compound must be determined by treating TA100 with a broad range of doses and observing the potential toxic effects on macrocolonies and microcolonies. If these tests are performed and expected data are obtained, then the results of the Ames test can be considered valid.

After validation of bacterial strains and selection of optimal sublethal doses, TRIETHYLENEGLYCOL DINITRATE was evaluated in the Ames Test. Criteria for a positive response include both a correlated dose-response over three dose concentrations and a revertant colony count at least two times (TA98, TA100) or three times (TA1535, TA1537, TA1538) the spontaneous revertant colony count (6). TRIETHYLENEGLYCOL DINITRATE did not induce the requisite dose response relationship or the increase in revertant colony counts necessary for a positive response. Thus, the results of this test indicate that TRIETHYLENEGLYCOL DINITRATE is not mutagenic when evaluated in the Ames Test.

TABLE 1: TOXICITY DETERMINATION FOR TEGON (TA044)

									
GLP	STUDY	NUMBER:	84048	16	JAN	85	PERFORMED	BY:	SANO

TA 100 REVERTANT PLATE COUNT

TEST COMPOUND CONCENTRATION	MEAN (1SD)	BACKGROUND LAWN*
NEGATIVE CONTROI, 5.0 µl/plate 1.0 µl/plate 0.2 µl/plate	95 (4.6) 99 (13.4) 96 (17.2) 107 (17.6)	NR NL NL NL
0.04 µl/plate 0.008 µl/plate 0.0016 µl/plate	85 (8.3) 110 (2.1) 97 (6.0)	NL NL NL

STRAIN VERIFICATION FOR TOXICITY DETERMINATION

TA 100*

HISTIDINE REQUIREMENT	NG
AMPICILLIN RESISTANCE	G
UV	NG
CRYSTAL VIOLET SENSITIVITY	NG (13mm)
STERILITY CONTROL	NG

STERILITY CONTROL FOR TOXICITY DETERMINATION

MATERIAL TESTED	OBSERVATION*
MINIMAL GLUCOSE AGAR PLATES	NG
TOP AGAR	NG
DILUENT	NG
NUTRIENT BROTH	NG
TEST COMPOUND (HIGHEST DOSE)	NG

^{*} NL=Normal Lawn, G=Growth, NG=No Growth, ST=Slight Toxicity NR=Not Recorded

TABLE 2: STRAIN VERIFICATION AND STERILITY TESTING FOR THE MUTAGENICITY DETERMINATION OF TEGDN (TA044)

GLP STUDY NUMBER: 84048 7-25 JAN 85 PERFORMED BY: SANO

STRAIN VERIFICATION

OBSERVATIONS*

STRAIN	HISTIDINE REQUIREMENT	AMPICILLIN RESISTANCE	UV REPAIR	CRYSTAL VIOLET (zone size)	CONTROL
TA98	NG	G	NG	NG (14mm)	NG
TA100	NG	G	NG	NG (10mm)	NG
TA1535	NG	NG	NG	NG (12mm)	NG
TA1537	NG	NG	NG	NG (13mm)	NG
TA1538	NG	NG	NG	NG (10mm)	NG

STERILITY CONTROL FOR MUTAGENICITY DETERMINATION

MATERIAL TESTED	OBSERVATION*
MINIMAL GLUCOSE AGAR PLATES TOP AGAR DILUENT WATER NUTRIENT BROTH TEST COMPOUND (HIGHEST DOSE) S-9	NG NG NG NG NG NG

^{*} NL=Normal Lawn, G=Growth, NG=No Growth, ST=Slight Toxicity

TABLE 3: MUTAGENICITY ASSAY FOR TEGDN (TA044)

REVERTANTS/PLATE (MEAN ±1SD)

COMPOUND	DOSE/PLATE	TA98	82	TA	TA100
		MITHOUT S-	ø		
	,				(
START NEG CONTROL	6 1 0.0		14.0	0	7.01
	2.0 49		ı	1017	±167.8
+ 0.44	51.0.00		1		ı
FINING		٧	v	42	10.
TAU44	11:		• •	7.8	ויכ ירט
TA044	7 11 1	r c		· C) a
TA044	0.2 µ1	.	7	107	. (
TA044		~	4.	9/	77
		۲	7	84	1.
# POKE	0 0016	4	<u>ر</u>	95	20
ENT NEC CONTROL	3	15	15.0	108	±7.9
	F	,	I		
		WITH S-9			
CHAPT NEC CONTROL	0.0 89	27	^'	98	15.
	2	~	366.	\sim	30.
2-25-4	2.0 Hg	444	13	533	±144.6
2 - 72 - 2	0	ω	29.5	0	19.
7704	=	~	1.5	σ	14.
TACAT	E [20	Ţ.	88	20.
PPOKE	۲c	24	ω,	72	8.4
**************************************		21	Α.	57	11.
4 0 0 4 F	0.008 11	26	-	0	30.
## OK E	•	27	ς.	2	14.
END NEG CONTROL	pn 0:0	24	+3.5	106	8.0
	n -				i

† MNNG = N-methyl-N'-nitro-N-nitrosoguanidine; 2-AA = 2-aminoanthracene; 2-AF aminofluorene; BP = benzo[a]pyrene.

MUTAGENICITY ASSAY FOR TEGDN (TA044) 3 (cont.): TABLE

REVERTANTS/PLATE (MEAN ±1SD)

COMPOUND	DOSE/PLATE	TA1535	TA1537	TAI	TA1538
		WITHOUT S-9	el.		
START NEG CONTROL	6n 0.0	13 ±1.5	8 ±2.0	23	±3.6
MINIGH	0	!	1	1	ı
MNNG+	0.0	~	1	•	ı
TA044	5 ml	3 ±4.	÷0.	30	±4.6
TA044		18 ±4.2	4 ±3.8	31	±10.4
TA044		1 ±3.	#	29	•
TA044	٠	5 ±2.	₽	44	•
TA044	0.008 µl	9 ±2.	Ħ.	35	∓6.0
TA044	•	7 ±5.	1 2.	35	•
END NEG CONTROL	0.0 µg	1 ±2.	Ħ.	29	1 5.9
		WITH S-9			
START NEG CONTROL	0.	13 ±2.5	#	28	•
2-AA+	2.0 µg	1	245 ±56.5	1286	02
2- A F†	0	1	1	559	2
BP↑	0	1	;	88	•
TA044	E	1 ±4.	#	25	•
TA044		2 +	Ħ.	19	•
TA044	•	7 ±3.	#3.	30	•
TA044	٥.	2 #	Ħ.	19	•
TA044	00.	7 ±4.	‡2·	27	•
4.	0.0016 µ1	18 ±4.4	5 ±1.2	30	±1.2
END NEG CONTROL	•	4	Ħ.	25	•

† MNNG = N-methyl-N'-nitro-N-nitrosoguanidine; 2-AA = 2-aminoanthracene; 2-AF = 2-aminofluorene; BP = benzo[a]pyrene.

Sano and Korte--10

CONCLUSION

TRIETHYLENEGLYCOL DINITRATE was evaluated for mutagenic potential in the Ames Test, in both the presence and absence of metabolic activation, and did not produce a positive mutagenic response under conditions of this study.

REFERENCES

- 1. Holleman JW, Ross RH, Carroll JW. Problem definition study on the health effects of diethyleneglycol dinitrate, triethyleneglycol dinitrate, and trimethylolethane trinitrate and their respective combustion products. Frederick, Maryland: US Army Medical Bioengineering Research and Development Laboratory, 1983, DTIC No. ADA 127846.
- 2. McCann JE, Choi E, Yamasaki E, Ames BN. Detection of carcinogens as mutagens in the Salmonella/Mammalian microsome test: Test of 300 chemicals. Proc Natl Acad Sci USA 1975;72:5135-5139.
- 3. Ames Salmonella/Mammalian Microsome Mutagenesis Test. LAIR Standard Operating Procedure OP-STX-1, Letterman Army Institute of Research, Presidio of San Francisco, California, 15 November 1983.
- 4. Ames BN, McCann J, Yamasaki E. Methods for detection of carcinogens and mutagens with Salmonella/Mammalian microsome mutagenicity test. Mutat Res 1975;31:347-364.
- 5. Vogel HJ, Bonner DM. Acetylornithinase of E. coli: Partial purification and some properties. J Biol Chem 1956;218:97-106.
- 6. Brusick D. Genetic Toxicology. In: Hayes AW, ed. Principles and Methods of Toxicology. New York: Raven Press, 1982: 223-272.

Sano and Korte--12

Appendix A.	Chemical Data13
Appendix B.	Individual Plate Scores

APPENDIX A: CHEMICAL DATA

Chemical Name: Ethanol, 2,2'-[1,2-ethanediylbis(oxy)] bis-,

dinitrate

Alternate Chemical Names: Triethyleneglycol dinitrate

(TEGDN), NOSET-A

Chemical Abstracts Service Registry No.: 111-22-8

LAIR Code No.: TA044

Chemical Structure:

O2N-O-CH2CH2-O-CH2CH2-O-CH2CH2-O-NO2

Molecular Formula: C6H12N2O8

Molecular Weight: 240

Physical State: Yellow oil

Density: (g/cm^3) : 1.32*

Source: Naval Ordnance Station, Indian Head, MD, 20640

Lot No.: 130-84

^{*} Holleman JW, Ross RH, Carroll JW. Problem definition study on the health effects of diethyleneglycol dinitrate, triethyleneglycol dinitrate and trimethylolethane trinitrate and their respective combustion products. Frederick, Maryland: US Army Medical Bioengineering Research and Development Laboratory, 1983, DTIC No. ADA 127846, p17.

APPENDIX A (cont.): CHEMICAL DATA

Analytical data: The compound chromatographed as a single peak (retention time 5.8 min) by HPLC analysis under the following conditions: column, Brownlee RP-18 (4.6 x 250 mm); solvent system, 30% water, 70% methanol; flow rate 0.9 ml/min, detection wavelength, 215 nm.† No impurities were detectable by NMR.‡ NMR (80 MHz, CDC13): 3.65 (S, 4H, -CH2-O-CH2CH2-O-CH2-), 3.72-3.84 (Complex multiplet, 4H, terminal methylene groups). IR (KBr): 2900, 1630, 1280, 1130, 1030, 910, 860 cm.\$

Stability: The compound was received as a 10% solution in ethanol. Periodic analysis of this solution by HPLC has shown no evidence of decomposition to date (4 months).† NMR analysis demonstrated that the neat compound is stable for at least 1 month.‡

[†] Wheeler, CR. Nitrocellulose-Nitroguanidine Projects. Laboratory Notebook #84-05-010.1, p26-30, 42-43. Letterman Army Institute of Research, Presidio of San Francisco, CA.

^{*} Wheeler, CR. Nitrocellulose-Nitroguanidine Projects. Laboratory Notebook #84-05-010.2, p63. Letterman Army Institute of Research, Presidio of San Francisco, CA.

^{§ &}lt;u>Ibid</u>. p64.

APPENDIX B: INDIVIDUAL PLATE SCORES

TRIETHYLENEGLYCOL DINITRATE (TA044) TOXICITY TEST TA100

0.04 µ1/p	78 94 82 normal lawn		
0.2 µ1/p	87 114 120 normal lawn	NEG CONTROL 90 98	not recorded
1.0 µ1/p	116 84 89 normal lawn	0.0016 µ1/p 97 91 103	normal lawn
5.0 µ1/p	114 89 93 normal lawn	0.008 µ1/p 111 112 108	normal lawn
DOSES	PLATE 1 PLATE 2 PLATE 3 background lawn	DOSES PLATE 1 PLATE 2 PLATE 3	background lawn

APPENDIX B (cont.): INDIVIDUAL PLATE SCORES

NEGATIVE CONTROLS FOR MUTAGENICITY TESTS TRIETHYLENEGLYCOL DINITRATE (TA044)

COMPOUND	DOSE	TA98	TA100	TA1535	TA1537	TA1538
WITHOUT S-9						
NEG CONTROL (START RUN)	0.0 µg/plate	20 12 16	91 77 97	15 12 13	8 10 6	20 22 22
NEG CONTROL (END RUN)	0.0 µg/plate	10 20 14	99 114 111	20 20 24	6 55 7	22 31 33
WITH S-9						
NEG CONTROL (START RUN)	0.0 µg/plate	28 25 29	74 81 103	13 11 16	0 0 B	29 24 24
NEG CONTROL (END RUN)	0.0 µg/plate	20 27 24	98 114 106	12 15 15	4 5 8	21 24 29

APPENDIX B (cont.): INDIVIDUAL PLATE SCORES

POSITIVE CONTROLS FOR MUTAGENICITY ASSAY TRIETHYLENEGLYCOL DINITRATE (TA044)

	DOSE	TA98	TA100	TA1535	TA15 37	TA1538
2-AA 2	2.0 µg/plate	1352 619 956	1600 1647 1657		288 181 266	1283 1185 1389
2-AF 2	2.0 µg/plate	370 600 361	369 586 643			600 613 464
ВР 2	2.0 µg/plate	206 148 186	500 532 496		4 4 4 4 8 8	93 86 86
MNNG 2	2.0 µg/plate		926 1211 915			
MNNG 2	20.0 µg/plate			2047 1619 1769		

APPENDIX B (cont.): INDIVIDUAL PLATE SCORES

TRIETHYLENEGLYCOL DINITRATE (TA044) MUTAGENICITY TESTS

WITHOUT S-9

TA044 5 μ1/plate 15 38 20 5 26 35 26 35 35 35 35 35 35 35 35 35 35 35 35 35 35 35 36 35 36 35 36 35 36	COMPOUND	DOSE	TA98	TA100	TA1535	TA1537	TA1538
1 μ1/plate 23 74 21 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	TA044	5 µl/plate	15 20 7	38 35 35		W W 4	
0.00 \(\mu1/\text{plate}\) 0.0016 \(\mu1/\text{plate}\) 25 \\ 106 \\ 25 \\ 107 \\ 25 \\ 107 \\ 25 \\ 107 \\ 26 \\ 27 \\ 90 \\ 27 \\ 90 \\ 28 \\ 90 \\ 21 \\ 34 \\ 71 \\ 110 \\ 25 \\ 72 \\ 20 \\ 72 \\ 20 \\ 72 \\ 20 \\ 72 \\ 20 \\ 72 \\ 20 \\ 72 \\ 20 \\ 73 \\ 74 \\ 44 \\ 25 \\ 72 \\ 25 \\ 72 \\ 27 \\	TA044	•	23 30 19	74 75 84		8 1 7	38 36 36
0.004 µ1/plate	TA044		18 16 25	00		887	3 3 3 3 3 8
0.008 µl/plate 20 91 20 6 3 28 90 21 4 2 34 71 16 3 4 0.0016 µl/plate 20 103 10 3 4 25 72 20 7 27 110 20 6 3	_A044		20 27 18	70 90 79	26 22 26	∞ 7 œ	
0.0016 µl/plate 20 103 10 3 4 25 72 20 7 27 110 20 6 3	_A044	ω	20 28 34	91 90 71	20 21 16	94 8	36 29 11
	TA044	16	20 25 27	103 72 110	10 20 20	e - 9	41 33 33

APPENDIX B (cont.): INDIVIDUAL PLATE SCOKES

TRIMETHYLOLETHANE TRINITRATE (TA044) MUTAGENICITY TESTS

WITH S-9

COMPOUND	DOSE	TA98	TA100	TA1535	TA1537	TA1538
TA044	5 µ1/plate	25	105	26	7	30
		23	79	17	· ഹ	20
		22	101	20	ഗ	2 <u>6</u>
TA044	1 µl/plate	20	29	19	7	7.0
		24	108	25	ഹ	· 60
		16	88	21	œ	11
TA044	0.2 µl/plate	21	62	14	α	96
		28	77	20	o	310
		23	92	18	7	33
TA044	0.04 µl/plate	21	64	<u>ر</u> بر	U	Č
	•	19	55.	13	n n	12
		22	46	6	۰ ۲	19
TA044	0.008 ul/plate	33	רר	7.2	u	Ġ
		25	108	18	ი	320
		19	117	20	° 7	25 29
TA044	0.0016 ul/plate	25	100	Ċ	•	į
		5 2	131	21	o 4	31
		30	104	13	9	29

Distribution List

Commander
US Army Biomedical Research and
Development Laboratory (27)
ATTN: SGRD-UBZ-C
Fort Detrick, Frederick, MD 21701-5010

Defense Technical Information Center (DTIC) (2)
ATTN: DTIC-DLA
Cameron Station
Alexandria, VA 22304-6145

US Army Medical Research and
Development Command (2)
ATTN: SGRD-RMI-S
Fort Detrick, Frederick, MD 21701-5012

Commandant Academy of Health Sciences, US Army ATTN: AHS-CDM Fort Sam Houston, TX 78234

Chief USAEHA Regional Division, West Fitzsimmons AMC Aurora, CO 80045

Chief USAEHA Regional Division, North Fort George G. Meade, MD 20755

Chief USAEHA Regional Division, South Bldg. 180 Fort McPherson, GA 30330

Commander
USA Health Services Command
ATTN: HSPA-P
Fort Sam Houston, TX 78234

Commandant
Academy of Health Sciences
United States Army
ATTN: Chief, Environmental
Quality Branch
Preventive Medicine Division
(HSHA-IPM)
Fort Sam Houston, TX 78234

Commander US Army Materiel Command ATTN: AMSCG 5001 Eisenhower Avenue Alexandria, VA 22333

Commander
US Army Environmental Hygiene
Agency
ATTN: Librarian, HSDH-AD-L
Aberdeen Proving Ground, MD 21010

Dean
School of Medicine
Uniformed Services University of the
Health Sciences
4301 Jones Bridge Road
Bethesda, MD 20014

Commander
US Army Materiel Command
ATTN: AMCEN-A
5001 Eisenhower Avenue
Alexandria, VA 22333

HQDA ATTN: DASG-PSP-E Falls Church, VA 22041-3258

HQDA ATTN: DAEN-RDM 20 Massachusetts, NW Washington, D.C. 20314